

Cellular structures using U_q -tilting modules

Or: centralizer algebras are fun!

Daniel Tubbenhauer

$$\begin{array}{ccccc} & & \Delta_q(\lambda) & & \\ & & \downarrow \iota^\lambda & \searrow g_i^\lambda & \\ M & \xrightarrow{\bar{r}_j^\lambda} & T_q(\lambda) & \xrightarrow{\bar{g}_i^\lambda} & N \\ & \searrow f_j^\lambda & \downarrow \pi^\lambda & & \\ & & \nabla_q(\lambda) & & \end{array}$$

Joint work with Henning Haahr Andersen and Catharina Stroppel

April 2015

The main theorem

Theorem

Let T be a $\mathbf{U}_q = \mathbf{U}_q(\mathfrak{g})$ -tilting module. Then $\text{End}_{\mathbf{U}_q}(T)$ is a cellular algebra.

I have to explain the words in red. But let us start with an example.

Example(Schur 1901)

Let $\mathbb{K}[S_d]$ be the symmetric group in d letters and let $\Delta_1(\omega_1)$ be the vector representation of $\mathbf{U}_1 = \mathbf{U}_1(\mathfrak{sl}_n)$. Take $T = \Delta_1(\omega_1)^{\otimes d}$, then

$$\Phi_{\text{SW}}: \mathbb{K}[S_d] \twoheadrightarrow \text{End}_{\mathbf{U}_1}(T) \quad \text{and} \quad \Phi_{\text{SW}}: \mathbb{K}[S_d] \xrightarrow{\cong} \text{End}_{\mathbf{U}_1}(T), \text{ if } n \geq d.$$

Since T is a \mathbf{U}_1 -tilting module, $\mathbb{K}[S_d]$ is cellular.

- 1 \mathbf{U}_q -tilting modules
 - \mathbf{U}_q and its representation theory
 - The category of \mathbf{U}_q -tilting modules
- 2 Cellularity of $\text{End}_{\mathbf{U}_q}(T)$
 - Cellular algebras
 - Cellularity and \mathbf{U}_q -tilting modules
- 3 The representation theory of $\text{End}_{\mathbf{U}_q}(T)$
 - Consequences of cellularity - \mathbf{U}_q -tilting view
 - Examples that fit into the picture

Quantum groups at roots of unity

Fix an arbitrary element $q \in \mathbb{K} - \{0\}$. Define

$$\mathbf{U}_q = \mathbf{U}_q(\mathfrak{g}) = \mathbf{U}_{\mathcal{A}} \otimes_{\mathcal{A}} \mathbb{K}.$$

Here $\mathbf{U}_{\mathcal{A}} = \mathbf{U}_{\mathcal{A}}(\mathfrak{g})$ is Lusztig's \mathcal{A} -form: the \mathcal{A} -subalgebra of $\mathbf{U}_v = \mathbf{U}_v(\mathfrak{g})$ generated by $K_i^{\pm 1}$, $E_i^{(j)}$ and $F_i^{(j)}$ for $i = 1, \dots, n-1$ and $j \in \mathbb{N}$.

Example

In the \mathfrak{sl}_2 case, the $\mathbb{Q}(v)$ -algebra $\mathbf{U}_v(\mathfrak{sl}_2)$ is generated by K, K^{-1} and E, F subject to some relations.

Let q be a complex, primitive third root of unity. $\mathbf{U}_q(\mathfrak{sl}_2)$ is generated by $K, K^{-1}, E, F, E^{(3)}$ and $F^{(3)}$ subject to some relations. Here $E^{(3)}, F^{(3)}$ are extra generators, since $E^3 = [3]!E^{(3)} = 0$ because of $[3] = 0$.

Weyl modules as building blocks

For each dominant \mathbf{U}_v -weight $\lambda \in X^+$ there is a simple \mathbf{U}_v -module $\Delta_v(\lambda)$ called Weyl module. Fact: the set $\{\Delta_v(\lambda) \mid \lambda \in X^+\}$ is a complete set of pairwise non-isomorphic, simple \mathbf{U}_v -modules (of type 1).

Example

For \mathfrak{sl}_2 we have $X^+ = \mathbb{Z}_{\geq 0}$. The Weyl module $\Delta_v(3)$ is

$$\begin{array}{ccccccc} \begin{array}{c} \curvearrowright \\ v^{-3} \\ \curvearrowleft \\ m_3 \end{array} & \begin{array}{c} \xrightarrow{[1]} \\ \xleftarrow{[3]} \end{array} & \begin{array}{c} \curvearrowright \\ v^{-1} \\ \curvearrowleft \\ m_2 \end{array} & \begin{array}{c} \xrightarrow{[2]} \\ \xleftarrow{[2]} \end{array} & \begin{array}{c} \curvearrowright \\ v^{+1} \\ \curvearrowleft \\ m_1 \end{array} & \begin{array}{c} \xrightarrow{[3]} \\ \xleftarrow{[1]} \end{array} & \begin{array}{c} \curvearrowright \\ v^{+3} \\ \curvearrowleft \\ m_0 \end{array} \end{array}$$

where E “acts to the right”, F “acts to the left” and K “acts as a loop”.

The category of finite dimensional \mathbf{U}_v -modules is semi-simple.

Weyl modules as building blocks?

Fact: the $\Delta_q(\lambda)$'s are no longer (semi-)simple in general. But they have unique simple heads $L_q(\lambda)$. Fact: the set $\{L_q(\lambda) \mid \lambda \in X^+\}$ is a complete set of pairwise non-isomorphic, simple \mathbf{U}_q -modules (of type 1).

Example

Let $\mathfrak{g} = \mathfrak{sl}_2$ and q be a complex, primitive third root of unity. $\Delta_q(3)$ is

$$\begin{array}{ccccccc} \begin{array}{c} \curvearrowright^{q^{-3}} \\ m_3 \end{array} & \begin{array}{c} \xrightarrow{+1} \\ \xleftarrow{0} \end{array} & \begin{array}{c} \curvearrowright^{q^{-1}} \\ m_2 \end{array} & \begin{array}{c} \xrightarrow{-1} \\ \xleftarrow{-1} \end{array} & \begin{array}{c} \curvearrowright^{q^{+1}} \\ m_1 \end{array} & \begin{array}{c} \xrightarrow{0} \\ \xleftarrow{+1} \end{array} & \begin{array}{c} \curvearrowright^{q^{+3}} \\ m_0 \end{array} \\ & & & & & \begin{array}{c} \xrightarrow{+1} \end{array} & \\ & & & & & \begin{array}{c} \xrightarrow{+1} \end{array} & \end{array}$$

The \mathbb{C} -span of $\{m_1, m_2\}$ is now stable under the action of $\mathbf{U}_q(\mathfrak{sl}_2)$: this is $L_q(1)$. The simple head is $L_q(3) \cong \Delta_q(3)/L_q(1)$ and is spanned by $\{m_0, m_3\}$.

The category of finite dimensional \mathbf{U}_q -modules is not semi-simple in general.

\mathbf{U}_q -tilting modules as building blocks?

Let $\Delta_q(\lambda)$ be a Weyl module and $\nabla_q(\lambda)$ its dual.

A \mathbf{U}_q -tilting module T is a \mathbf{U}_q -module with a Δ_q -filtration and a ∇_q -filtration:

$$\begin{aligned} T &= M_0 \supset M_1 \supset \cdots \supset M_{k'} \supset \cdots \supset M_{k-1} \supset M_k = 0, \\ 0 &= N_0 \subset N_1 \subset \cdots \subset N_{k'} \subset \cdots \subset N_{k-1} \subset N_k = T, \end{aligned}$$

such that $M_{k'}/M_{k'+1}$ is some $\Delta_q(\lambda)$ and $N_{k'+1}/N_{k'}$ is some $\nabla_q(\lambda)$.

Example

All \mathbf{U}_v -modules are \mathbf{U}_v -tilting modules.

For our favorite example $q^3 = 1 \in \mathbb{C}$ and $\mathfrak{g} = \mathfrak{sl}_2$: $\Delta_q(i)$ is a \mathbf{U}_q -tilting module iff $i = 0, 1$ or $i \equiv -1 \pmod{3}$.

\mathbf{U}_q -tilting modules as building blocks.

The category of \mathbf{U}_q -tilting modules \mathcal{T} has some nice properties:

- \mathcal{T} is closed under finite tensor products.
- The indecomposables $T_q(\lambda)$ of \mathcal{T} are parametrized by $\lambda \in X^+$. They have λ as their maximal weight and contain $\Delta_q(\lambda)$ with multiplicity 1. We have

$$\Delta_q(\lambda) \xrightarrow{\iota^\lambda} T_q(\lambda) \xrightarrow{\pi^\lambda} \nabla_q(\lambda).$$

Example

The vector representation $\Delta_q(1)$ is a $\mathbf{U}_q(\mathfrak{sl}_2)$ -tilting module. Thus, $T = \Delta_q(1)^{\otimes d}$ is. Then $T_q(d)$ is the indecomposable summand of T containing $\Delta_q(d)$.

Example

$\Delta_q(\lambda)$ is a \mathbf{U}_q -tilting module for minuscule λ . Thus, tensor products of these are.

The Ext-vanishing

We have for all $\lambda, \mu \in X^+$ that

$$\mathrm{Ext}_{\mathbf{U}_q}^i(\Delta_q(\lambda), \nabla_q(\mu)) \cong \begin{cases} \mathbb{K}c^\lambda, & \text{if } i = 0 \text{ and } \lambda = \mu, \\ 0, & \text{else,} \end{cases}$$

where $c^\lambda: \Delta_q(\lambda) \rightarrow \nabla_q(\lambda)$ is the \mathbf{U}_q -homomorphism that sends head to socle.

Assume that M has a Δ_q -filtration and N has a ∇_q -filtration.

- We have $\dim(\mathrm{Hom}_{\mathbf{U}_q}(M, \nabla_q(\lambda))) = (M : \Delta_q(\lambda))$.
- We have $\dim(\mathrm{Hom}_{\mathbf{U}_q}(\Delta_q(\lambda), N)) = (N : \nabla_q(\lambda))$.

\mathbf{U}_q -tilting modules as building blocks!

$$T \in \mathcal{T} \quad \text{iff} \quad \text{Ext}_{\mathbf{U}_q}^1(T, \nabla_q(\lambda)) = 0 = \text{Ext}_{\mathbf{U}_q}^1(\Delta_q(\lambda), T) \quad \text{for all } \lambda \in X^+.$$

In particular, if M has a Δ_q - and N has a ∇_q -filtration:

$$\begin{array}{ccccc}
 & & \Delta_q(\lambda) & & \\
 & & \downarrow \iota^\lambda & \searrow g_i^\lambda & \\
 M & \xrightarrow{\bar{f}_j^\lambda} & T_q(\lambda) & \xrightarrow{\bar{g}_i^\lambda} & N \\
 & \searrow f_j^\lambda & \downarrow \pi^\lambda & & \\
 & & \nabla_q(\lambda) & &
 \end{array}$$

In words: any \mathbf{U}_q -homomorphism $g: \Delta_q(\lambda) \rightarrow N$ extends to an \mathbf{U}_q -homomorphism $\bar{g}: T_q(\lambda) \rightarrow N$ whereas any \mathbf{U}_q -homomorphism $f: M \rightarrow \nabla_q(\lambda)$ factors through $T_q(\lambda)$ via $\bar{f}: M \rightarrow T_q(\lambda)$.

Consequence of the discussion before:

$$\dim(\text{End}_{\mathbf{U}_q}(T)) = \sum_{\lambda \in X^+} (T : \Delta_q(\lambda))^2 = \sum_{\lambda \in X^+} (T : \nabla_q(\lambda))^2.$$

Take $T = \Delta_q(\lambda)^{\otimes d}$. If $\lambda \in X^+$ is minuscule as a \mathbf{U}_q -weight, then $\Delta_q(\lambda)$ is always \mathbf{U}_q -tilting and $\dim(\text{End}_{\mathbf{U}_q}(T))$ is independent of \mathbb{K} and q , since $\Delta_q(\lambda)$ has a character independent of \mathbb{K} and of q .

Example

By quantum Schur-Weyl, we see that

$$\Phi_{q\text{SW}} : \mathcal{H}_d(q) \twoheadrightarrow \text{End}_{\mathbf{U}_q}(T) \quad \text{and} \quad \Phi_{q\text{SW}} : \mathcal{H}_d(q) \xrightarrow{\cong} \text{End}_{\mathbf{U}_q}(T), \text{ if } n \geq d.$$

Thus, $\dim(\mathcal{H}_d(q))$ independent of \mathbb{K} and q .

Exempli gratia (Temperley-Lieb without diagrams)

Let us consider our favorite case again. From the construction of $T_q(3)$:

$$\Delta_q(3) \hookrightarrow T_q(3) \twoheadrightarrow \Delta_q(1).$$

We compute:

$$T_v = \Delta_v(1) \otimes \Delta_v(1) \otimes \Delta_v(1) \cong \Delta_v(3) \oplus \Delta_v(1) \oplus \Delta_v(1),$$

whereas

$$T_q = \Delta_q(1) \otimes \Delta_q(1) \otimes \Delta_q(1) \cong T_q(3) \oplus T_q(1).$$

In particular, $\dim(\text{End}_{\mathbf{U}_v(\mathfrak{sl}_2)}(T_v)) = \dim(\text{End}_{\mathbf{U}_q(\mathfrak{sl}_2)}(T_q)) = 1^2 + 2^2 = 5$.

Note that $\text{End}_{\mathbf{U}_q(\mathfrak{sl}_2)}(\Delta_q(1)^{\otimes d})$ is the Temperley-Lieb algebra $\mathcal{TL}_d(\delta)$.

Definition (Graham-Lehrer 1996)

A \mathbb{K} -algebra A is cellular if it has a basis

$$\{c_{ij}^\lambda \mid \lambda \in \mathcal{P}, i, j \in \mathcal{I}\},$$

where (\mathcal{P}, \leq) is a finite poset and \mathcal{I}^λ is a finite set, such that

- 1 The map $i: A \rightarrow A, c_{ij}^\lambda \rightarrow c_{ji}^\lambda$ is an anti-isomorphism.
- 2 We have (for friend of higher order)

$$ac_{ij}^\lambda = \sum_{k \in \mathcal{I}^\lambda} r_{ik}(a)c_{kj}^\lambda + \text{friends.}$$

Note that the scalars $r_{ik}(a)$ do not depend on j . Thus, we think of the basis elements as having “independent bottom and top parts”.

Prototype of a cellular basis

Example(Specht 1935, Murphy 1995)

\mathcal{P} = Young diagrams λ , \mathcal{I}^λ = standard tableaux i, j .

$$c_{ij}^\lambda = \begin{array}{c} \begin{array}{|c|c|c|} \hline \dots & & \\ \hline \end{array} \\ \begin{array}{|c|c|c|} \hline P^*(i) & & \\ \hline \end{array} \text{ permutation} \\ \begin{array}{|c|c|c|} \hline e(\lambda) & & \\ \hline \end{array} \text{ idempotent} \\ \begin{array}{|c|c|c|} \hline \dots & & \\ \hline \end{array} \\ \begin{array}{|c|c|c|} \hline P(j) & & \\ \hline \end{array} \text{ permutation} \\ \begin{array}{|c|c|c|} \hline \dots & & \\ \hline \end{array} \end{array}$$

Form $S^\lambda = \{c_j^\lambda\}$ with formal c_j^λ and action given by the $r_{ik}(a)$. The set

$$\{D^\lambda = S^\lambda / \text{Rad}(S^\lambda) \mid \lambda \in \mathcal{P}_0\}$$

forms a complete set of pairwise non-isomorphic, simple $\mathbb{K}[S_d]$ -modules.

Theorem(Graham-Lehrer 1996)

This works in general for cellular algebras.

And for $\text{End}_{\mathbf{U}_q}(T)$?

Let M have a Δ_q - and N have ∇_q -filtration. Consider $\mathcal{I}^\lambda = \{1, \dots, (N : \nabla_q(\lambda))\}$ and $\mathcal{J}^\lambda = \{1, \dots, (M : \Delta_q(\lambda))\}$. By Ext-vanishing, we have diagrams

$$\begin{array}{ccc}
 M & \xrightarrow{\bar{f}_j^\lambda} & T_q(\lambda) \\
 & \searrow f_j^\lambda & \downarrow \pi^\lambda \\
 & & \nabla_q(\lambda)
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 \Delta_q(\lambda) & & \\
 \downarrow \iota^\lambda & \searrow g_i^\lambda & \\
 T_q(\lambda) & \xrightarrow{\bar{g}_i^\lambda} & N
 \end{array}$$

Take any bases $F^\lambda = \{f_j^\lambda : M \rightarrow \nabla_q(\lambda) \mid j \in \mathcal{J}^\lambda\}$ of $\text{Hom}_{\mathbf{U}_q}(M, \nabla_q(\lambda))$ and $G^\lambda = \{g_i^\lambda : \Delta_q(\lambda) \rightarrow N \mid i \in \mathcal{I}^\lambda\}$ of $\text{Hom}_{\mathbf{U}_q}(\Delta_q(\lambda), N)$. Set

$$c_{ij}^\lambda = \bar{g}_i^\lambda \circ \bar{f}_j^\lambda \in \text{Hom}_{\mathbf{U}_q}(M, N)$$

for each $\lambda \in X^+$ and all $i \in \mathcal{I}^\lambda, j \in \mathcal{J}^\lambda$.


$\text{End}_{\mathbf{U}_q}(T)$ is prototypical cellular

Cell datum:

- $(\mathcal{P}, \leq) = (\{\lambda \in X^+ \mid (T : \nabla_q(\lambda)) = (T : \Delta_q(\lambda)) \neq 0\}, \leq_X)$.
- $\mathcal{I}^\lambda = \{1, \dots, (T : \nabla_q(\lambda))\} = \{1, \dots, (T : \Delta_q(\lambda))\} = \mathcal{J}^\lambda$ for each $\lambda \in \mathcal{P}$.
- \mathbb{K} -linear anti-involution $i: \text{End}_{\mathbf{U}_q}(T) \rightarrow \text{End}_{\mathbf{U}_q}(T), \phi \mapsto \mathcal{D}(\phi)$.
- Note that $\mathcal{D}(\Delta_q(\lambda)) \cong \nabla_q(\lambda)$ and $\mathcal{D}(\nabla_q(\lambda)) \cong \Delta_q(\lambda)$.
- Cellular basis $\{c_{ij}^\lambda \mid \lambda \in \mathcal{P}, i, j \in \mathcal{I}^\lambda\}$.

Theorem


This gives a cellular datum on $\text{End}_{\mathbf{U}_q}(T)$ for any \mathbf{U}_q -tilting module T .



Exempli gratia (generic Temperley-Lieb)

Take $\mathbb{K} = \mathbb{C}$ and $T = \Delta_v(1)^{\otimes 3} \cong \Delta_v(3) \oplus \Delta'_v(1) \oplus \Delta''_v(1)$. Then $\mathcal{P} = \{1, 3\}$.


We have $\mathcal{I}^1 = \{1, 2\}$ and $\mathcal{I}^3 = \{1\}$. Thus, we have a basis



Exempli gratia (roots of unity Temperley-Lieb)

Take $T = \Delta_q(1)^{\otimes 3} \cong T_q(3) \oplus T_q(1)$. Then $\mathcal{P} = \{1, 3\}$.

We have $\mathcal{I}^1 = \{1, 2\}$ and $\mathcal{I}^3 = \{1\}$. Consider $1 \in \mathcal{I}^1$ as indexing the factor $\Delta_q(1)$ of $T_q(1)$ and $2 \in \mathcal{I}^1$ the factor $\Delta_q(1)$ of $T_q(3)$. Thus, we have a basis



Cellular pairing and simple $\text{End}_{\mathbf{U}_q}(T)$ -modules

Let T be a \mathbf{U}_q -tilting module. For $\lambda \in \mathcal{P}$ define ϑ^λ via

$$i(h) \circ g = \vartheta^\lambda(g, h)c^\lambda, \quad g, h \in C(\lambda) = \text{Hom}_{\mathbf{U}_q}(\Delta_q(\lambda), T).$$

Define $\mathcal{P}_0 = \{\lambda \in \mathcal{P} \mid \vartheta^\lambda \neq 0\}$ and $\text{Rad}(\lambda) = \{g \in C(\lambda) \mid \vartheta^\lambda(g, C(\lambda)) = 0\}$.

Theorem(Graham-Lehrer reinterpreted)

The set

$$\{L(\lambda) = C(\lambda)/\text{Rad}(\lambda) \mid \lambda \in \mathcal{P}_0\}$$

is a complete set of pairwise non-isomorphic, simple $\text{End}_{\mathbf{U}_q}(T)$ -modules.

$\lambda \in \mathcal{P}_0$ iff $T_q(\lambda)$ is a summand of T . Moreover,

$$\dim(L(\lambda)) = m_\lambda, \quad T \cong \bigoplus_{\lambda \in X^+} T_q(\lambda)^{\oplus m_\lambda}.$$

Exempli gratia (Temperley-Lieb again)

Because $T_v \cong \Delta_v(3) \oplus \Delta_v(1) \oplus \Delta_v(1)$ and $T_q \cong T_q(3) \oplus T_q(1)$ we see that $\mathcal{P}_0 = \{1, 3\}$ in both cases.

In the generic case:

$$C(3) = L(3) = \{g_1^3: \Delta_v(3) \rightarrow T_v\}, \quad C(1) = L(1) = \{g_j^1: \Delta_v(1) \rightarrow T_v \mid j = 1, 2\},$$
$$\dim(L(3)) = 1 \quad \text{and} \quad \dim(L(1)) = 2.$$

In the non-semisimple case:

$$C(3) = L(3) = \{g_1^3: \Delta_q(3) \rightarrow T_q\}, \quad C(1) = \{g_j^1: \Delta_q(1) \rightarrow T_q \mid j = 1, 2\},$$
$$\dim(L(3)) = 1 \quad \text{and} \quad \dim(L(1)) = 1.$$

An alternative semi-simplicity criterion

Theorem (Graham-Lehrer 1996)

Let A be a cellular algebra with cell modules $C(\lambda)$ and simple modules $L(\lambda)$.

$$A \text{ is semi-simple} \Leftrightarrow C(\lambda) = L(\lambda) \text{ for all } \lambda \in \mathcal{P}_0.$$

We can prove an alternative statement in our framework.

Theorem

The algebra $\text{End}_{\mathbf{U}_q}(T)$ is semi-simple iff T is a semi-simple \mathbf{U}_q -module.

Corollary

The algebra $\text{End}_{\mathbf{U}_q}(T)$ is semi-simple iff T has only simple Weyl factors.

Exempli gratia (Temperley-Lieb yet again)

Because $T_v \cong \Delta_v(3) \oplus \Delta_v(1) \oplus \Delta_v(1)$, and $\Delta_v(3)$ and $\Delta_v(1)$ are simple Weyl factors, we see that $\text{End}_{\mathbf{U}_v(\mathfrak{sl}_2)}(T_v)$ is semi-simple.

T_q has a Weyl factor of the form $\Delta_q(3)$. This is a non-simple Weyl factor and thus, $\text{End}_{\mathbf{U}_q(\mathfrak{sl}_2)}(T_q)$ is non semi-simple.

Similarly: $\mathcal{TL}_d(\delta) \cong \text{End}_{\mathbf{U}_q(\mathfrak{sl}_2)}(\Delta_q(1)^{\otimes d})$ with $\delta \neq 0$ is semi-simple iff q is not a root of unity in \mathbb{K} or $d < \text{ord}(q^2)$.

A unified approach to cellularity - part 1

Note that our approach generalizes, for example to the infinite dimensional world: the following list is just the tip of the iceberg.

The following algebras fit in our set-up as well:

- The Iwahori-Hecke algebra of type A , by Schur-Weyl duality:

$$\Phi_{q\text{SW}}: \mathcal{H}_d(q) \rightarrow \text{End}_{\mathbf{U}_q}(T) \quad \text{and} \quad \Phi_{q\text{SW}}: \mathcal{H}_d(q) \xrightarrow{\cong} \text{End}_{\mathbf{U}_q}(T), \text{ if } n \geq d.$$

This includes $\mathbb{K}[S_d]$ for $\text{char}(\mathbb{K}) = p > 0$.

- \mathfrak{sl}_2 -related algebras like Temperley-Lieb $\mathcal{TL}_d(\delta)$.
- Spider algebras $\text{End}_{\mathbf{U}_q(\mathfrak{sl}_n)}(\Delta_q(\omega_{i_1}) \otimes \cdots \otimes \Delta_q(\omega_{i_d}))$.

A unified approach to cellularity - part 2

- Take $\mathfrak{g} = \mathfrak{gl}_{m_1} \oplus \cdots \oplus \mathfrak{gl}_{m_r}$ with $m_1 + \cdots + m_r = m$ and let V be the vector representation of $\mathbf{U}_1(\mathfrak{gl}_m)$ restricted to $\mathbf{U}_1 = \mathbf{U}_1(\mathfrak{g})$. Use $T = V^{\otimes d}$ and

$$\Phi_{\text{cl}}: \mathbb{C}[\mathbb{Z}/r\mathbb{Z}\lambda S_d] \rightarrow \text{End}_{\mathbf{U}_1}(T) \quad \text{and} \quad \Phi_{\text{cl}}: \mathbb{C}[\mathbb{Z}/r\mathbb{Z}\lambda S_d] \xrightarrow{\cong} \text{End}_{\mathbf{U}_1}(T), \quad \text{if } m \geq d.$$

This gives the cyclotomic analogon of the first point above.

- Let $\mathbf{U}_q = \mathbf{U}_q(\mathfrak{g})$. We get in the quantized case

$$\Phi_{q\text{cl}}: \mathcal{H}_{d,r}(q) \rightarrow \text{End}_{\mathbf{U}_q}(T) \quad \text{and} \quad \Phi_{q\text{cl}}: \mathcal{H}_{d,r}(q) \xrightarrow{\cong} \text{End}_{\mathbf{U}_q}(T), \quad \text{if } m \geq d,$$

where $\mathcal{H}_{d,r}(q)$ is the Ariki-Koike algebra.

A unified approach to cellularity - part 3

- Let $T = \Delta_q(\omega_1)^{\otimes d}$. Let $\mathfrak{g} = \mathfrak{o}_{2n}$, $\mathfrak{g} = \mathfrak{o}_{2n+1}$ or $\mathfrak{g} = \mathfrak{sp}_{2n}$ (depending on δ).

$$\Phi_{\text{Br}}: \mathcal{B}_d(\delta) \twoheadrightarrow \text{End}_{\mathbf{U}_1}(T) \quad \text{and} \quad \Phi_{\text{Br}}: \mathcal{B}_d(\delta) \xrightarrow{\cong} \text{End}_{\mathbf{U}_1}(T), \text{ if } 2n > d,$$

where $\mathcal{B}_d(\delta)$ is the Brauer algebra in d strands.

- Let $\mathbf{U}_q = \mathbf{U}_q(\mathfrak{gl}_n)$ and $T = \Delta_q(\omega_1)^{\otimes r} \otimes (\Delta_q(\omega_1)^{\otimes s})^*$:

$$\Phi_{\text{wBr}}: \mathcal{B}_{r,s}^n([n]) \twoheadrightarrow \text{End}_{\mathbf{U}_q}(T) \quad \text{and} \quad \Phi_{\text{wBr}}: \mathcal{B}_{r,s}^n([n]) \xrightarrow{\cong} \text{End}_{\mathbf{U}_q}(T), \text{ if } n \geq r+s,$$

where $\mathcal{B}_{r,s}^n([n])$ the so-called quantized walled Brauer algebra.

- Quantizing the Brauer case: taking $q \in \mathbb{K} - \{0, \pm 1\}$, \mathfrak{g} , and T as above: the algebra $\text{End}_{\mathbf{U}_q}(T)$ is a quotient of the Birman-Murakami-Wenzl algebra $\mathcal{BMW}_d(\delta)$ and taking $n \geq d$ recovers $\mathcal{BMW}_d(\delta)$.

There is still **much** to do...

Thanks for your attention!